

# Growth Factor Data Sheet

GoldBio growth factors are manufactured for **RESEARCH USE ONLY** and cannot be sold for human consumption!

FGF7/KGF1 is a member of the Fibroblast Growth Factor family of proteins. In the FGF family, FGF7 is most closely related to FGF10 (KGF2), FGF3 and FGF22 (see additional information). Proteins of this family play a central role during prenatal development and postnatal growth and regeneration of a variety of tissues, by promoting cellular proliferation and differentiation. FGF7/KGF1 is mitogen factor specific for epithelial cells and keratinocytes and signals through Fibroblast Growth Factor Receptor-2b (FGFR-2b) and plays an important role in kidney and lung development, angiogenesis and wound healing.

<b>Catalog Number</b>	<b>1340-07</b>
<b>Product Name</b>	<b>FGF7/KGF1, Murine</b> Recombinant Murine Fibroblast Growth Factor 7 FGF7 KGF1 Keratinocyte Growth Factor 1
<b>Source</b>	<i>Escherichia coli</i>
<b>MW</b>	~18.9 kDa (164 amino acids)
<b>Sequence</b>	CNDMSPEQTA TSVNCSSEPER HTRSYDMEG GDIRVRLFC RTQWYLRIK RGKVKGTEQEM KNSYNIMEIR TVAVGIVAIAK GVESEYLLAM NKEGKLYAKK ECNEDCNFKE LILENHYNLY ASAKWTHSGG EMFVALNQKG IPVKGKTKK EQKTAHFLPM AIT
<b>Accession Number</b>	<a href="#">P36363</a>
<b>Purity</b>	>97% by SDS-PAGE and HPLC analyses
<b>Biological Activity</b>	Fully biologically active when compared to standard. The ED <sub>50</sub> as determined by thymidine uptake assay using FGF-receptors transfected BaF3 cells is less than 10 ng/ml, corresponding to a specific activity of >1.0 × 10 <sup>5</sup> IU/mg.
<b>Formulation</b>	Sterile filtered white lyophilized powder. Purified and tested for use in cell culture.
<b>Storage/Handling</b>	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage. The reconstituted sample can be apportioned into working aliquots and stored at -80 °C for up to 6 months. Avoid repeated freeze/thaw cycles.
<b>Reconstitution</b>	The sample should be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a siliconized tube using PBS that contains a 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Reconstituted solutions are stable for up to one week at 2-8°C. Stock solutions should be aliquoted and stored at -80°C. Further dilutions should be made in appropriate buffered solutions containing BSA or serum.