

Growth Factor Data Sheet

GoldBio growth factors are manufactured for **RESEARCH USE ONLY** and cannot be sold for human consumption!

Interleukin-13 (IL13) is expressed by the IL13 gene and secreted by many cell types, especially type 2 helper T-cells (Th2). The high solution form of IL13 has been reported to be a monomer with two internal disulfide bonds that contribute to its bundled four F-helix configuration. Targeted deletion of IL13 in mice resulted in impaired Th2 cell development and suggested an important role for IL13 in the expulsion of gastrointestinal parasites. IL13 exerts anti-inflammatory effects on monocytes and macrophages, and it inhibits the expression of inflammatory cytokines such as IL1 β , TNF- α , IL6 and IL8. IL13 has also been shown to enhance B-cell proliferation and induce isotype switching, which results in an increased production of IgE. Mature rhesus IL13 shares 95%, 59% and 60% amino acid sequence identity with human, mouse, and rat IL13, respectively.

Catalog Number	1210-13
Product Name	IL13, Rhesus macaque Recombinant Rhesus macaque Interleukin-13 IL13 Interleukin 13
Source	<i>Escherichia coli</i>
MW	~12.6 kDa (114 amino acids)
Sequence	SPSPVPRSTA LKELIEELVN ITQNQKAPLC NGSMVWSINL TAGVYCAALE SLINVSGCSA IEKTQRMLNG FCPHKVSAGQ FSSLRVRDTK IEVAQFVKDL LVHLKFLFRE GRFN
Accession Number	Click here to enter text.
Purity	>98% by SDS-PAGE and HPLC analyses
Biological Activity	Q864V6
Formulation	Sterile filtered white lyophilized powder. Purified and tested for use in cell culture.
Storage/Handling	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage. The reconstituted sample can be apportioned into working aliquots and stored at -80 °C for up to 6 months. Avoid repeated freeze/thaw cycles.
Reconstitution	The sample should be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a siliconized tube using PBS that contains a 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Reconstituted solutions are stable for up to one week at 2-8°C. Stock solutions should be aliquoted and stored at -80°C. Further dilutions should be made in appropriate buffered solutions containing BSA or serum.