

Growth Factor Data Sheet

GoldBio growth factors are manufactured for **RESEARCH USE ONLY** and cannot be sold for human consumption!

Tumor necrosis factor (or TNF- α), also called cachectin, is produced by neutrophils, activated lymphocytes, macrophages, NK-cells, LAK-cells, astrocyte endothelial cells, smooth muscle cells and some transformed cells. TNF occurs as a secreted, soluble form and as a membrane-anchored form, both of which are biologically active. The naturally-occurring form of TNF is glycosylated, but nonglycosylated recombinant TNF has comparable biological activity. The biologically active, native form of TNF is reportedly a trimer. Human and murine TNF has approximately 79% homology at the amino acid level and there is cross-reactivity between the two species. Two types of receptors for TNF, p55 and p75, have been described and virtually all cell types studied show the presence of one or both of these receptor types.

Catalog Number	1120-03
Product Name	TNF-α, Human Recombinant Human Tumor Necrosis Factor TNF- α , TNF-alpha Cachectin, Cachexin
Source	<i>Escherichia coli</i>
MW	~17.5 kDa (158 amino acid)
Sequence	MVRSSRTPS DKPVAHVVAN PQAEGQLQWL NRRANALLAN GVELRDNQLV VPSEGLYLIY SQVLFKGQGC PSTHVLLTHT ISRIAVSYQT KVNLLSAIKS PCQRETPEGA EAKPWYEPIY LGGVFLQLEKG DRLSAEINRP DYLDFAESGQ VYFGIIAL
Accession Number	P01375
Purity	>98% by SDS-PAGE and HPLC analyses
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by cytotoxicity assay using murine L929 cells is less than 0.05 ng/ml in the presence of actinomycin D.
Formulation	Sterile filtered white lyophilized powder. Purified and tested for use in cell culture.
Storage/Handling	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage. The reconstituted sample can be apportioned into working aliquots and stored at -80 °C for up to 6 months. Avoid repeated freeze/thaw cycles.
Reconstitution	The sample should be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a siliconized tube using PBS that contains a 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Reconstituted solutions are stable for up to one week at 2-8°C. Stock solutions should be aliquoted and stored at -80°C. Further dilutions should be made in appropriate buffered solutions containing BSA or serum.