

Growth Factor Data Sheet

GoldBio growth factors are manufactured for RESEARCH USE ONLY and cannot be sold for human consumption!

IL22 belongs to IL10 family of regulatory cytokines, which includes IL10, IL19, IL20, IL22, IL24 and IL26. Members of this family share partial homology in their amino acid sequences, but are dissimilar in their biological functions. Produced by T-lymphocytes and dendritic cells, IL10 contributes to the inflammatory response *in vivo*, and signals through CRF2-4 and IL22. Along with IL17, it is rapidly produced by splenic LTi-like cells and can be also produced by Th17 cells. It is likely that IL22 plays a role in the coordinated response of both the adaptive and innate immune systems.

Catalog Number	1510-22
Product Name	IL22, Rat Recombinant Rat Interleukin-22 IL22 Interleukin 22 Interleukin 10-Related T-cell-derived Inducible Factor (ILTIF)
Source	<i>Escherichia coli</i>
MW	~16.6 kDa (146 amino acids)
Sequence	LPINSQCKLE AANFQQPYIV NRTFMLAKEA SLADNNTDVR LIGEELFRGV KAKDQCYLKMK QVLNFTLEDV LLPQSDRFQP YMQEVPVFLT KLSIHLSPCH ISGDDQNIQK NVRQLKETVQ KLGESGEIKA IGELDLLFMS LRNACV
Accession Number	Q198B4
Purity	>97% by SDS-PAGE and HPLC analyses
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by inducing IL-10 secretion of human COLO 205 cells is less than 0.5 ng/ml, corresponding to a specific activity of >2.0 × 10 ⁶ IU/mg.
Formulation	Sterile filtered white lyophilized powder. Purified and tested for use in cell culture.
Storage/Handling	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage. The reconstituted sample can be apportioned into working aliquots and stored at -80 °C for up to 6 months. Avoid repeated freeze/thaw cycles.
Reconstitution	The sample should be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a siliconized tube using PBS that contains a 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Reconstituted solutions are stable for up to one week at 2-8°C. Stock solutions should be aliquoted and stored at -80°C. Further dilutions should be made in appropriate buffered solutions containing BSA or serum.