

Growth Factor Data Sheet

GoldBio growth factors are manufactured for **RESEARCH USE ONLY** and cannot be sold for human consumption!

Interferon gamma is the only type II interferon found in humans and binds to the heterodimeric IFNGR that consists of two chains, IFNGR1 and IFNGR2. This interaction activates the JAK-STAT pathway. IFNG is produced primarily by T-lymphocytes and natural killer (NK) cells. It is critical for innate and adaptive immunity against viral and intracellular bacterial infections. This is partly due to its ability to inhibit viral replication directly, but also due to its immunostimulatory and immunomodulatory effects. IFNG induces the production of cytokines and upregulates the expression of class I and II MHC antigens, Fc receptors, and leukocyte adhesion molecules. It modulates macrophage effector functions, influences isotype switching, and potentiates the secretion of immunoglobulins by B cells. Additionally, IFNG augments TH1 cell expansion and may be required for TH1 cell differentiation.

Catalog Number	1160-06
Product Name	IFN-gamma, Human Recombinant Human Interferon Gamma IFNG IFG IFN, Immune (IFI)
Source	<i>Escherichia coli</i>
MW	~16.9 kDa (144 amino acids)
Sequence	MQDPYVKEAE NLKRYFNAGH SDVADNGTLF LGILKNWKEE SDRKIMQSQI VSFYFKLFKN FKDDQSIQKS VETIKEDMNV KFFNSNKKKR DDFEKLTNYS VTDLNVQRKA IHELIQVMAE LSPAAGTGKR KRSQMLFRGR RASQ
Accession Number	P01579
Purity	>98% by SDS-PAGE and HPLC analyses
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as measured in anti-viral assay using human HeLa cells infected with encephalomyocarditis (EMC) virus is 0.15-0.80 ng/ml.
Formulation	Sterile filtered white lyophilized powder. Purified and tested for use in cell culture.
Storage/Handling	The lyophilized sample is stable at 2-8°C, but should be kept desiccated at -20°C for long term storage. The reconstituted sample can be apportioned into working aliquots and stored at -80 °C for up to 6 months for maximal stability. Avoid repeated freeze/thaw cycles.
Reconstitution	The sample should be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a siliconized tube using PBS that contains a 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Reconstituted solutions are stable for up to one week at 2-8°C. Stock solutions should be aliquoted and stored at -80°C. Further dilutions should be made in appropriate buffered solutions containing BSA or serum.