

Growth Factor Data Sheet

GoldBio growth factors are manufactured for **RESEARCH USE ONLY** and cannot be sold for human consumption!

CCL28 is a chemokine containing six conserved cysteine residues, two of which are adjacent. It is expressed constitutively in epithelial cells at mucosal sites such as salivary glands and colon and expression is upregulated by proinflammatory stimuli such as IL1 and n-butyrate. CCL28 is chemotactic for eosinophils, resting T cells, and IgA antibody-secreting cells (ASCs). It is important for regulating IgA ASC accumulation in mammary glands and consequently the transfer of IgA antibodies from mother to infant. It also plays a role in tumor growth and angiogenesis. CCL28 is a ligand for CCR10 and CCR3.

Catalog Number	2140-28
Product Name	CCL28, Human Recombinant Human Chemokine (C-C motif) Ligand 28 (CCL28) Mucosae-Associated Eptihelial Chemokine (MEC) Small Inducible Cytokine Subfamily A Member 28 (SCYA28)
Source	<i>Escherichia coli</i>
MW	~12.4 kDa (108 amino acids)
Sequence	SEAILPIASS CTEVSHHIS RLLERVNMC RIQRADGDCD LAAVILHVKR RRICVSPHNN TVKQWMKVQA AKKNGKGNVC HRKKHHGKRN SNRAHQGKHE TYGHKTPY
Accession Number	Click here to enter text.
Purity	>97% by SDS-PAGE and HPLC analyses
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human lymphocytes is in a concentration range of 1.0-10.0 ng/ml.
Formulation	Sterile filtered white lyophilized powder. Purified and tested for use in cell culture.
Storage/Handling	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage. The reconstituted sample can be apportioned into working aliquots and stored at -80 °C for up to 6 months. Avoid repeated freeze/thaw cycles.
Reconstitution	The sample should be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a siliconized tube using PBS that contains a 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Reconstituted solutions are stable for up to one week at 2-8°C. Stock solutions should be aliquoted and stored at -80°C. Further dilutions should be made in appropriate buffered solutions containing BSA or serum.